ORAL MICROBIOLOGY AND IMMUNOLOGY

Detection of oral bacteria in cardiovascular specimens

Nakano K, Nemoto H, Nomura R, Inaba H, Yoshioka H, Taniguchi K, Amano A, Ooshima T. Detection of oral bacteria in cardiovascular specimens.

Oral Microbiol Immunol 2009: 24: 64–68. © 2009 The Authors. Journal compilation. © 2009 Blackwell Munksgaard.

Background/aims: Oral bacteria, including cariogenic and periodontal pathogens, are thought to be etiological factors in the development of cardiovascular diseases. To define this relationship, we analyzed the distribution of oral bacterial species in cardiovascular specimens.

Method: Following acceptance into the study, 203 consecutive patients were analyzed, from whom 82 aortic valve specimens, 35 mitral valve specimens, and 86 aortic aneurysmal wall specimens, of which 16 contained aneurysmal thrombus tissues, were obtained. In addition, a total of 58 dental plaque specimens were collected from the same group of patients who underwent heart valve replacement or removal of aortic aneurysms. Bacterial DNA was extracted from both cardiovascular tissues and dental plaque in those cases and then species-specific polymerase chain reaction assays were used to analyze the occurrences of six oral streptococcal and six periodontal bacterial species.

Results: *Streptococcus mutans* was the most frequently detected species in the cardiovascular specimens, followed by *Aggregatibacter actinomycetemcomitans*. As for dental plaque specimens from patients who underwent cardiovascular operations, most of the tested periodontitis-related species as well as oral streptococci were detected at high frequencies. Furthermore, the positive rate of *S. mutans* in cardiovascular specimens from patients whose dental plaque specimens were also positive for *S. mutans* was 78%, which was significantly higher than any other tested species when the same analysis was performed.

Conclusion: Our results suggest that specific oral bacterial species, such as *S. mutans* and *A. actinomycetemcomitans*, are related to bacteremia and may be etiologic factors for the development of cardiovascular diseases.

K. Nakano¹, H. Nemoto¹, R. Nomura¹, H. Inaba², H. Yoshioka³, K. Taniguchi⁴, A. Amano², T. Ooshima¹

¹Department of Pediatric Dentistry, Division of Oral Infection and Disease Control, and ²Department of Oral Frontier Biology, Division of Oral Biology and Disease Control, Osaka University Graduate School of Dentistry, Suita, Osaka, Japan, ³Department of Dentistry and Oral Surgery, and ⁴Department of Cardiovascular Surgery, Osaka Rosai Hospital, Sakai, Osaka, Japan

Key words: aortic aneurysm; heart valve; molecular approach; polymerase chain reaction; *Streptococcus mutans*

Takashi Ooshima, Department of Pediatric Dentistry, Division of Oral Infection and Disease Control, Osaka University Graduate School of Dentistry, 1-8 Yamadaoka, Suita, Osaka 565-0871, Japan Tel.: +81 6 6879 2961; fax: +81 6 6879 2965; e-mail: ooshima@dent.osaka-u.ac.jp

Accepted for publication August 11, 2008

Dental caries and marginal periodontitis are two of the major conditions generally encountered in the field of dentistry, with both considered to be caused by pathogenic oral bacterial species (18, 19). It is becoming more apparent that oral microbes comprise a complex community, and that oral health or disease depends on the interface between the host and microbial community as a whole (6). Oral bacteria are also known to cause systemic diseases, such as infective endocarditis, which has been shown to be associated with oral streptococci (15). Furthermore, the association of periodontitis with cardiovascular diseases has recently become an issue of great concern.

In the field of cardiovascular surgery, heart valve disease and aortic aneurysms are major diseases. Recent populationbased surveys in the USA found that the national prevalence of valve diseases was approximately 2.5% in adults (17). These valve diseases mostly develop in aortic or mitral valves, and are diagnosed as regurgitation, stenosis, or both (2). On the other hand, aortic aneurysms are defined as permanent localized dilation of the aorta with a diameter at least 1.5 times greater than the expected normal diameter of the given aortic segment, and are mainly classified into thoracic aortic aneurysms (TAAs), abdominal aortic aneurysms (AAAs) and thoracoabdominal aortic aneurysms (TAAAs), based on the location of the diseased aorta (5). The estimated prevalence of aortic aneurysms in the USA was recently reported to be 1–2% of the adult population (10).

In our previous report, we noted that oral streptococci and periodontitis-related bacteria were detected in cardiovascular lesions extirpated from 62 patients using polymerase chain reaction (PCR) assays (16). However, the number of these cases studied was not adequate to provide conclusive results, while no other systematic large-scale study demonstrating the presence of oral bacterial species in cardiovascular tissues has been conducted. In the present study, we collected heart valve and aneurysm clinical specimens from 203 patients with various clinical diagnoses, and analyzed the prevalence of six oral streptococcal and six periodontitis-related bacterial species in them, to define the relationship between oral bacteria and cardiovascular diseases. In addition, we determined if the occurrences of these bacteria were related to cardiovascular pathologic conditions and diseased sites. Dental plaque samples were also collected from some of the patients, and used to analyze the relationships between oral and cardiovascular bacterial profiles.

Materials and methods Cardiovascular and dental plaque specimens

The study procedures were approved by the Ethics Committee of Osaka Rosai Hospital and informed consent was obtained from each subject. Two hundred and three consecutive cases were analyzed, from which 82 aortic valve specimens, 35 mitral valve specimens, and 86 aortic aneurysmal wall specimens, 16 of which contained aneurysmal thrombus tissue, were obtained at the Department of Cardiovascular Surgery, Osaka Rosai Hospital, from December 2004 to November 2007. Table 1 summarizes the characteristics of the cases, such as number of patients, diagnosis, gender, and age. In addition, supragingival and subgingival plaque samples from some of these patients, referred for oral care before their cardiovascular operations, were taken from the mesial and buccal subgingival sites of all teeth with sterile Gracey curettes at the Department of Dentistry and Oral Surgery of the same hospital. A total of 58 dental plaque specimens were collected from 29 patients who underwent heart valve replacement and 29 patients who underwent removal of aortic aneurysms. The cardiovascular and dental plaque specimens were stored in sterile saline after collection with sterile instruments, then preserved at -20°C until analysis, as described previously (16).

Detection of 12 oral bacterial species

Six oral streptococcal species (*Streptococcus mutans*, *Streptococcus sobrinus*, *Streptococcus sobrinus*, *Streptococcus salivarius*, *Streptococcus sanguinis*, *Streptococcus oralis*, and *Streptococcus gordonii*), as well as six periodontitis-related species (*Porphyromonas gingivalis*, *Prevotella intermedia*, *Treponema denticola*, *Tannerella forsythia*, *Aggregatibacter actinomycetemcomitans*, *Campylobacter rectus*) were detected with PCR using species-specific sets of primers, as described previously (1, 4, 9, 22, 23). The primers used and other details are shown in Table 2.

Statistical analysis

Statistical analyses were carried out using the computational software package STAT-VIEW 5.0 (SAS Institue Inc., Cary, NC). Fisher's protected least-significant difference test was utilized to compare the detection frequency of each species.

Results

Distribution of oral bacterial species in cardiovascular specimens

The detection rates of all tested species were classified according to clinical diagnosis based on pathological and anatomical conditions, as shown in Fig. 1. A majority of the specimens (80% of heart valves, 90% of aortic aneurysms) were found to contain more than one species. There were no significant relationships between bacterial occurrences and clinical classification of the specimens. Regardless of diagnosis, S. mutans was the most frequently detected species, with detection rates in the heart valve and aneurysm wall specimens of 42.7% and 62.8%, respectively, while A. actinomycetemcomitans was the second most often detected, in 35.0% and 30.2%, respectively. As for the other species, S. sanguinis, P. gingivalis, and T. denticola were detected at frequencies ranging from 15 to 20%, each of which was significantly lower than S. mutans (P < 0.05), while the detection rates of S. sobrinus, S. salivarius, S. oralis, S. gordonii, P. intermedia, T. forsythia, and C. rectus were extremely low.

Detection of oral bacterial species in dental plaque and cardiovascular specimens from the same patients

The bacterial profiles of the dental plaque and cardiovascular specimens collected from the same patients were compared (Fig. 2). Periodontitis-related species were often detected in dental plaque, at rates of more than 70% for *C. rectus*, *T. forsythia*, and *T. denticola*, while the occurrences of *P. gingivalis* and *A. actinomycetemcomitans* were 50–60%. In contrast, *P. intermedia*, the least often occurring periodontitis-related species, was detected

Table 1. Summary of cases analyzed in the present study

	Number of c	ases			
Clinical diagnosis	Male	Female	Total	Age^1 [mean \pm SD (years)]	
Aortic regurgitation (AR)	27	4	31	66.8 ± 9.1 (36–84)	
Aortic stenosis (AS)	15	17	32	$69.1 \pm 8.1 \ (49-81)$	
Aortic stenosis and regurgitation (ASR)	9	10	19	67.8 ± 11.3 (46-87)	
Total aortic valves	51	31	82	$67.9 \pm 9.2 \ (36-87)$	
Mitral regurgitation (MR)	6	17	23	64.0 ± 8.4 (49–78)	
Mitral stenosis (MS)	3	2	5	$66.8 \pm 2.9 (63 - 71)$	
Mitral stenosis and regurgitation (MSR)	3	4	7	69.6 ± 9.7 (58–84)	
Total mitral valves	12	23	35	$65.5 \pm 8.3 \ (49-84)$	
Abdominal aortic aneurysm (AAA)	39	8	47	$70.6 \pm 7.1 \ (56-84)$	
Thoracic aortic aneurysm (TAA)	18	10	28	69.4 ± 8.0 (53–78)	
Thoracoabdominal aortic aneurysm (TAAA)	8	3	11	73.8 ± 5.7 (64–82)	
Total aneurysms	65	21	86	$69.9 \pm 9.8 (53 - 84)$	

¹Parentheses indicate the range of age distribution.

66 Nakano et al.

Table 2.	List of	polymerase	chain	reaction	primers	used	in	the	present	study
		/								

Purpose	Sequence (5' to 3')	Size (bp)	Reference
Detection of oral streptococci			
S. mutans	GGC ACC ACA ACA TTG GGA AGC TCA GTT	433	(4)
	GGA ATG GCC GCT AAG TCA ACA GGA T		
S. sobrinus	GAT GAT TTG GCT CAG GAT CAA TCC TC	328	(4)
	ACT GAG CCA GTA GTA GAC TTG GCA ACT		
S. salivarius	GTG TTG CCA CAT CTT CAC TCG CTT CGG	544	(4)
	CGT TGA TGT GCT TGA AAG GGC ACC ATT		
S. sanguinis	GGA TAG TGG CTC GGG CAG CCA GTT	313	(4)
-	GAA CAG TTG CTG GAC TTG CTT GTC		
S. oralis	TCC CGG TCA GCA AAC TCC AGC C	374	(4)
	GCA ACC TTT GGA TTT GCA AC		
S. gordonii	CTA TGC GGA TGA TGC TAA TCA AGT G	440	(4)
-	GGA GTC GCT ATA ATC TTG TCA GAA A		
Detection of periodontitis-related species			
Porphyromonas gingivalis	TGT AGA TGA CTG ATG GTG AAA ACC	197	(22)
	ACG TCA TCC CCA CCT TCC TC		
Prevotella intermedia	TTT GTT GGG GAG TAA AGC GGG	575	(1)
	TCA ACA TCT CTG TAT CCT GCG T		
Treponema denticola	TAA TAC CGA ATG TGC TCA TTT ACA T	311	(23)
	TCA AAG AAG CAT TCC CTC TTC TTC TTA		
Tannerella forsythia	GCG TAT GTA ACC TGC CCG CA	641	(1)
	TGC TTC AGT GTC CAG TTA TAC CT		
Aggregatibacter actinomycetemcomitans	CTA GGT ATT GCG AAA CAA TTT G	262	(9)
	CCT GAA ATT AAG CTG GTA ATC		
Campylobacter rectus	TTT CGG AGC GTA AAC TCC TTT TC	598	(1)
	TTT CTG CAA GCA GAC ACT CTT		



Fig. 1. Detection of oral bacterial species in cardiovascular specimens. The detection rates of the tested bacterial species in 117 heart valve specimens (A and B), 86 aortic aneurysmal wall specimens (C), and 102 aortic aneurysm specimens (D) were determined by polymerase chain reaction using species-specific sets of primers. Abbreviations: SMU, *Streptococcus mutans*; SSO, *Streptococcus sobrinus*; SSL, *Streptococcus salivarius*; SSA, *Streptococcus sanguinis*; SOR, *Streptococcus oralis*; SGO, *Streptococcus gordonii*; Pg, *Porphyromonas gingivalis*; Pi, *Prevotella intermedia*; Td, *Treponema denticola*; Tf, *Tannerella forsythia*; Aa, *Aggregatibacter actinomycetemcomitans*; Cr, *Campylobacter rectus*.

at a frequency of approximately 20%. Oral streptococcal species, including *S. mutans*, *S. sanguinis*, and *S. oralis*, were also often

detected in the dental plaque samples at rates of more than 60%, while the occurrences of *S. sobrinus*, *S. salivarius*, and *S. gordonii* were limited to approximately 10–15%. As shown in Fig. 2, there was a slight tendency for species prevalent in



Fig. 2. Detection of oral bacterial species in patients from whom both dental plaque and cardiovascular specimens were collected (n = 58). The detection rates of the tested bacterial species in dental plaque and cardiovascular specimens from patients who underwent heart valve replacement (A; n = 29) and aortic aneurysm removal (B; n = 29) procedures were determined by polymerase chain reaction using species-specific sets of primers. Abbreviations as in Fig. 1.

dental plaque to be detected in cardiovascular tissues, though this was not statistically significant.

Next, cardiovascular specimens from patients whose dental plaque specimens were positive for each species were further analyzed for bacterial occurrence (Table 3). We suspected that some oral bacteria were able to efficiently enter and spread in the bloodstream, which may be an etiologic event for the development of cardiovascular diseases. *S. mutans* had a detection rate of 78.0% in the cardiovascular specimens, which was significantly greater than any of the other tested species (P < 0.01). In addition, the detection rates of *A. actinomycetemcomitans* and *T. denticola* were 46.4% and 27.7%, respectively. In contrast, *P. gingivalis* was detected in only 5.6% of the cases and the occurrences of *S. oralis, S. gordonii*, and *T. forsythia* were also low (<3%).

Discussion

Dental treatment is known to be the major causative agent for bacteremia and procedures that generally cause bleeding, such as tooth extraction, periodontal surgery, scaling and root planing, are considered to frequently lead to its development (21). Furthermore, bacteremia has been reported after daily routine oral hygiene procedures including tooth brushing (0-26%) and dental flossing (20-58%), and even from chewing (17-51%) (20). In fact, a majority of the present specimens (approximately 80% of the heart valves, 90% of the aortic aneurysms) were positive for at least one of the tested oral bacterial species. In addition, there were no significant differences in detection rates for each species between aortic and mitral valves. as well as between valves associated with stenosis and regurgitation in the present study. These findings indicate that oral bacteria frequently enter the bloodstream and may invade cardiovascular tissues following dental treatments and after the performance of routines of daily life.

Recently, an association between periodontitis and cardiovascular diseases has been suggested, based on reports that bacterial DNA corresponding to periodontitis-related species was detected in cardiovascular tissues using PCR. Although several studies have described the detection of oral bacteria in cardiovascular specimens, few have performed simultadetection of gram-positive neous (oral streptococcal) and gram-negative (periodontitis-related) species. To our knowledge, this is the first report of both gram-positive and gram-negative bacterial profiles analyzed using cardiovascular and dental plaque specimens collected from a large number of subjects. In the present study, we focused on the detection of six oral streptococcal and six periodontal bacterial species, of which three periodontitis-related species and two oral streptococcal species were detected in more than 10% of cardiovascular specimens, with the detection rate of S. mutans being high (42.7% of the heart valve and 62.8% of aortic aneurysm specimens). In addition, S. mutans was detected in cardiovascular specimens with a significantly higher frequency in those subjects whose dental plaque specimens also showed a positive reaction for this species, as compared to the other species. A previous study that analyzed bacterial species in blood from child subjects using a blood culture technique also showed that the incidence of S. mutans tended to be higher than that of other species (12). Therefore, S. mutans likely enters the bloodstream more readily than the others, which may indicate it as a possible etiological factor for cardiovascular disease.

T.1.1. 2	Detection neter of	· 1	1			f		1	1 1				41 · · ·	
Ianie 1	Detection rates of	orai	nacteria in	carmovascui	ar specimens	from 1	nanents	whose	dental	niadiii	e snecimens	nossessed	the sam	- snecles
raore 5.	Detection fates of	orui	oucteria m	curatovascai	ar speemiens	nom	patiento	11000	aontai	praga	e speennens	possessed	the built	species
					-	-	-				-	-		-

Species ¹	Patients with	heart valve	replaceme	nt	Patients with	aortic valve	removal		Total patients with cardiovascular operations			
	Detection rate ² (%)	P-values ³			Detection	P-values ³			Detection	P-values ³		
		vs. SMU	vs. Aa	vs. Td	rate ² (%)	vs. SMU	vs. Aa	vs. Td	rate ² (%)	vs. SMU	vs. Aa	vs. Td
SMU	17/26 (65.4)	_	NS	NS	22/24 (91.7)	_	< 0.001	< 0.001	39/50 (78.0)	_	0.006	< 0.001
SSO	0/2 (0)	NS	NS	NS	1/4 (25.0)	0.012	NS	NS	1/6 (16.7)	0.006	NS	NS
SSL	0/5 (0)	0.012	0.045	NS	1/4 (25.0)	0.012	NS	NS	1/9 (11.1)	< 0.001	NS	NS
SSA	2/22 (9.1)	< 0.001	0.006	0.016	5/24 (20.8)	< 0.001	NS	NS	7/46 (15.2)	< 0.001	0.006	NS
SOR	0/20 (0)	< 0.001	< 0.001	0.001	1/17 (5.9)	< 0.001	NS	NS	1/37 (2.7)	< 0.001	0.001	0.002
SGO	0/2(0)	NS	NS	NS	0/3 (0)	0.003	NS	NS	0/5 (0)	0.001	NS	NS
Pg	1/18 (5.6)	< 0.001	0.004	0.011	1/18 (5.6)	< 0.001	NS	NS	2/36 (5.6)	< 0.001	< 0.001	0.010
Pi	1/9 (11.1)	0.007	0.040	NS	1/7 (14.3)	< 0.001	NS	NS	2/16 (12.5)	< 0.001	0.045	NS
Td	9/21 (42.9)	NS	NS	_	4/26 (15.4)	< 0.001	NS	_	13/47 (27.7)	< 0.001	NS	_
Tf	1/28 (3.6)	< 0.001	< 0.001	0.001	0/23 (0)	< 0.001	0.005	NS	1/51 (2.0)	< 0.001	< 0.001	< 0.001
Aa	8/14 (57.1)	NS	_	NS	5/14 (35.7)	0.001	_	NS	13/28 (46.4)	0.006	_	NS
Cr	6/26 (23.1)	0.005	0.043	NS	0/28 (0)	< 0.001	0.002	0.047	6/54 (11.1)	< 0.001	< 0.001	0.043

¹Bacterial species are abbreviated as follows: SMU, *Streptococcus mutans*; SSO, *Streptococcus sobrinus*; SSL, *Streptococcus salivarius*; SSA, *Streptococcus sanguinis*; SOR, *Streptococcus oralis*; SGO, *Streptococcus gordonii*; Pg, *Porphyromonas gingivalis*; Pi, *Prevotella intermedia*; Td, *Treponema denticola*; Tf, *Tannerella forsythia*; Aa, *Aggregatibacter actinomycetemcomitans*; Cr, *Campylobacter rectus*.

²The detection rate is expressed as the ratio of cardiovascular specimens positive for the species in all individuals whose dental plaque specimens were positive for the same species.

³NS indicates not statistically significant (i.e. P > 0.05).

68 Nakano et al.

In a previous study, oral cariogenic and periodontal conditions were epidemiologically evaluated using clinical and radiographic examinations of patients with cardiac valve disease, and the results were used to analyze the relationships of oral conditions with cardiovascular diagnostic variations (8). It appeared that aortic valve disorders developed significantly more often than mitral site disorders in patients with severely infected oral cavities. Although no clinical records with regard to the oral cavity were available for the present subjects, there were no significant differences found for the detection rates of the tested oral streptococcal and periodontal bacterial species between the aortic and mitral valve specimens. In addition, the rates of detection for oral bacterial species between valves extracted under the diagnosis of stenosis and regurgitation were not statistically different. Therefore, bacterial colonization in blood is not likely affected by the anatomical and pathological characteristics of the disease.

The association of oral pathogens with cardiovascular diseases has recently received attention, with atherosclerotic lesions caused by P. gingivalis infection being widely investigated. Among those results, systemic challenge by P. gingivalis was reported to accelerate atherogenic plaque formation in an apolipoprotein E-deficient murine model (11). On the other hand, several previous reports identified multiple oral bacterial species in human atheromatous plaque specimens (3, 7, 13, 14). In the present study, multiple oral bacterial species were detected in aortic aneurysm specimens and the detection rate for each species was similar to that of those species found in the heart valve specimens. Therefore, it is possible to speculate that the species detected in the aortic aneurysm specimens were related to bacteremia. However, additional investigations are required, because the species identified in these specimens might also be relevant for the pathogenesis of aortic aneurysm formation.

We collected dental plaque specimens from only about 30% of the present subjects with cardiovascular disease because the patients were originally examined at the Department of Cardiovascular Surgery and did not have an opportunity to also visit the Department of Dentistry and Oral Surgery, unless there was a specific problem in the oral cavity. Nevertheless, we frequently detected oral bacterial species in the cardiovascular specimens, which might be correlated with the development of cardiovascular diseases. Therefore, it is important for cardiovascular physicians and surgeons to understand the importance of oral health care and of routine dental examinations before surgery, while additional analyses focused on the direct involvement of oral bacteria in cardiovascular diseases may provide important findings.

Acknowledgments

This study was supported by the 21st Century COE program entitled 'Origination of Frontier BioDentistry' at the Osaka University Graduate School of Dentistry supported by the Ministry of Education, Culture, Sports, Science and Technology of Japan Grants-in-Aid for Scientific Research (A) 19209063 and (B) 16390605 from the Japan Society for Promotion of Science, and Grants-in-Aid for Young Scientists (A) 18689050 and (B)19791572 from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

- Ashimoto A, Chen C, Bakker I, Slots J. Polymerase chain reaction detection of 8 putative periodontal pathogens in subgingival plaque of gingivitis and advanced periodontitis lesions. Oral Microbiol Immunol 1996: 11: 266–273.
- Braunwald E. Valvular heart diseases. In: Braunwald E, Zipes DP, Libby P, eds. Heart disease, 6th edn. Philadelphia: W.B. Saunders, 2001: 1643–1722.
- Haraszthy VI, Zambon JJ, Trevisan M, Zeid M, Genco RJ. Identification of periodontal pathogens in atheromatous plaques. J Periodontol 2000: 71: 1554–1560.
- Hoshino T, Kawaguchi M, Shimizu N, Hoshino N, Ooshima T, Fujiwara T. PCR detection and identification of oral streptococci in saliva samples using *gtf* genes. Diagn Microbiol Infect Dis 2004: 48: 195– 199.
- Isselbacher EM. Diseases of the aorta. In: Braunwald E, Zipes DP, Libby P, eds. Heart disease, 6th edn. Philadelphia: W.B. Saunders, 2001: 1422–1456.
- Jenkinson HF, Lamont RJ. Oral microbial communities in sickness and in health. Trends Microbiol 2005: 13: 585–595.
- Kozarov E, Sweier D, Shelburne C, Progulske-Fox A, Lopatin D. Detection of bacterial DNA in atheromatous plaques by quantitative PCR. Microb Infect 2006: 8: 687–693.
- 8. Krennmair G, Auer J, Krainhöfner M, Piehslinger E. Odontogenic infection

sources in patients scheduled for cardiac valve replacement. Oral Health Prev Dent 2007: **2**: 153–159.

- Kuboniwa M, Amano A, Kimura KR et al. Quantitative detection of periodontal pathogens using real-time polymerase chain reaction with TaqMan probes. Oral Microbiol Immunol 2004: 19: 168–176.
- Kuivaniemi H, Platsoucas CD, Tilson MD 3rd. Aortic aneurysms: an immune disease with a strong genetic component. Circulation 2008: 117: 242–252.
- Li L, Messas E, Batista EL Jr, Levine RA, Amar S. *Porphyromonas gingivalis* infection accelerates the progression of atherosclerosis in a heterozygous apolipoprotein E-deficient murine model. Circulation 2002: 105: 861–867.
- Lockhart PB, Brennan MT, Kent ML, Norton HJ, Weinrib DA. Impact of amoxicillin prophylaxis on the incidence, nature, and duration of bacteremia in children after intubation and dental procedures. Circulation 2004: 109: 2878–2884.
- Marques da Silva R, Caugant DA, Eribe ER et al. Bacterial diversity in aortic aneurysms determined by 16S ribosomal RNA gene analysis. J Vasc Surg 2006: 44: 1055–1060.
- Meurman JH, Sanz M, Janket SJ. Oral health, atherosclerosis, and cardiovascular disease. Crit Rev Oral Biol Med 2004: 15: 403–413.
- Moreillon P, Que YA. Infective endocarditis. Lancet 2004: 363: 139–149.
- Nakano K, Inaba H, Nomura R et al. Detection of cariogenic *Streptococcus mutans* in extirpated heart valve and atheromatous plaque specimens. J Clin Microbiol 2006: 44: 3313–3317.
- Nkomo VT, Gardin JM, Skelton TN, Gottdiener JS, Scott CG, Enriquez-Sarano M. Burden of valvular heart diseases: a population-based study. Lancet 2006: 368: 1005–1011.
- Pihlstrom BL, Michalowicz BS, Johnson NW. Periodontal diseases. Lancet 2005: 366: 1809–1820.
- 19. Selwitz RH, Ismail AI, Pitts NB. Dental caries. Lancet 2007: 369: 51–59.
- Seymour RA, Lowry R, Whitworth JM, Martin MV. Infective endocarditis, dentistry and antibiotic prophylaxis; time for a rethink? Br Dent J 2000: 189: 610–616.
- Tomás Carmona I, Diz Dios P, Scully C. Efficacy of antibiotic prophylactic regimens for the prevention of bacterial endocarditis of oral origin. J Dent Res 2007: 86: 1142– 1159.
- 22. Tran SD, Rudney JD. Multiplex PCR using conserved and species-specific 16S rRNA gene primers for simultaneous detection of *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis*. J Clin Microbol 1996: 34: 2674–2678.
- Watanabe K, Frommel TO. Porphyromonas gingivalis, Actinobacillus actinomycetemcomitans and Treponema denticola detection in oral plaque samples using the polymerase chain reaction. J Clin Periodontol 1996: 23: 212–219.